



<https://www.coelomycetes.org>: Databank that contributes for the classification, identification and nomenclature of Coelomycetes

Huanraluek N¹, Padaruth OD¹, Jayawardena RS^{1,2,3}, Li WJ¹¹, Hongsanan S⁴, Jeewon R⁵, Maharachchikumbura SSN⁶, Senanayake IC^{1,2,3}, Phukhamsakda C^{7,8}, Wijayawardene NN⁹, Shivas RG¹⁰, Goonasekara ID^{1,2}, Norphanphoun C^{1,2}, Pem D^{1,2}, Calabon MS^{1,2}, Bhunjun CS^{1,2} and Hyde KD^{1,2,3*}

¹Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

²School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

³Innovative Institute for Plant Health, Zhongkai University of Agriculture and Engineering, Haizhu District, Guangzhou 510225, People's Republic of China

⁴Shenzhen Key Laboratory of Microbial Genetic Engineering, College of Life Sciences and Oceanography, Shenzhen University, Shenzhen 518060, P.R. China

⁵Department of Health Sciences, Faculty of Medicine and Health Science, University of Mauritius, Reduit, Mauritius

⁶School of Life Science and Technology, Center for Informational Biology, University of Electronic Science and Technology of China, Chengdu 611731, P.R. China

⁷Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi Jilin Agricultural University, Changchun City, Jilin Province, P.R. China, 130118

⁸College of Plant Protection, Jilin Agricultural University, Changchun, Jilin, P.R. China, 130118

⁹Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University, Qujing, Yunnan 655011, P.R. China

¹⁰Institute for Life Sciences and the Environment, Research and Innovation Division University of Southern Queensland, Toowoomba, Queensland, 4350, Australia

¹¹Yunnan Minzu University, Kunming 650504, People's Republic of China

Huanraluek N, Padaruth OD, Jayawardena RS, Li WJ, Hongsanan S, Jeewon R, Maharachchikumbura SSN, Senanayake IC, Phukhamsakda C, Wijayawardene NN, Shivas RG, Goonasekara ID, Norphanphoun C, Pem D, Calabon MS, Bhunjun CS and Hyde KD 2021 – <https://www.coelomycetes.org>: Databank that contributes for the classification, identification, and nomenclature of Coelomycetes. Asian Journal of Mycology 4(2), 114–122, Doi 10.5943/ajom/4/2/8

Abstract

Over the years, there have been continuous updates in the taxonomy of coelomycetes. In particular, traditional classification has been revised and updated based on extensive collections supplemented with morphological examinations and phylogenetic analyses. Most of the data available are scattered in the literature. In an attempt to ease the work of mycologists and bring all these data together, a website has been created, available at <https://www.coelomycetes.org>. Information provided on the webpage is in line with the current ranking and classification of taxa and will include photoplates, drawings, phylogenetic trees, notes and descriptions.

Keywords – asexual fungi – conidiogenesis – curators – database – website

Introduction

Coelomycetes is an artificial group known as imperfect fungi. It was introduced by Grove (1919) to accommodate the asexual morphs of fungi such as *Phomopsis*, *Phloeospora* and

Phyllosticta in *Ascomycetes*. Furthermore, members have been reported to be also related to asexually reproducing basidiomycetous fungi such as *Basidiopycnis*, *Chaetospermum* and *Fibulocoela* (Nag Raj 1981, Wijayawardene et al. 2016, Liu et al. 2019, Li et al. 2020). Asexual fungi are diverse and can survive in a variety of ecological environments (Hyde et al. 2020). They can be found in soil, leaf litter, organic or manufactured sources, from fresh or saline water as well as on lichens (Jayawardena et al. 2019, Li et al. 2020). Furthermore, these imperfect fungi can live as parasites on plants, humans, vertebrates or insects (Sutton 1980).

Coelomycetes was rejected as a formal taxonomic rank and later was used as a collective term in *Ascomycota* and *Basidiomycota* (Nag Raj 1993, Li et al. 2020). Several studies have accepted accept conidiogenesis as a primary taxonomic criterion in asexual fungi (Nag Raj 1993). Wijayawardene et al. (2016) and Li et al. (2020) agree with Kendrick (2000) that coelomycetes is a term used for convenience. However, the classification of coelomycetes excluded hyphomycetes and agonomycetes because genera in coelomycetes are separated based on their conidiomata structure in combination with conidiogenesis and conidia characteristics (Sutton 1980, Nag Raj 1993, Wijayawardene et al. 2016, Gruyter et al. 2017, Li et al. 2020, Wijayawardene et al. 2020). Furthermore, conidiomata differed in terms of morphology as they have conidiophores, conidiogenous cells and conidial while ascomatal structures are available in sexual fungi. In addition, coelomycete genera can produce conidia from an acervulus, pycnidial, sporodochial-like and synnema-like those bear conidia (Sutton 1980, Nag Raj 1993, Wijayawardene et al. 2016, Li et al. 2020, Wijayawardene et al. 2020).

Conidiomata have distinct differentiation of tissue types that are assumed to be either plastic or stable. During conidia ontogeny, the cell wall either grows or is left behind for conidium initiation (Sutton 1980, Nag Raj 1993). Hughes (1953) gave an experimental description of conidiogenesis to explain the production of conidia. Sutton (1980) also used conidiogenesis as a primary characteristic to identify coelomycetes and later, Nag Raj (1993) illustrated 142 genera and over 200 appendages bearing coelomycetes with descriptions. Based on DNA sequence phylogenetic analyses, most of the species accepted by both Sutton (1980) and Nag Raj (1993) were reclassified in different orders, classes, or genera (Li et al. 2020). For example, *Tiarosporella abeietis*, *T. parca* and *T. pseudotsugae* were transferred to *Darkera* (Leotiomyces) while *T. tritici* was placed under *Eutiarosporella* (*Botryosphaerales*) using morphological characters and phylogenetic results (Li et al. 2020). These show that, when using phylogenetic analyses, it is easier to place many species in their natural classification. However, it is common to have contradictions between phenotypic and genotypic data as pointed out by Li et al. (2020). The latter noted that the phylogenetic analyses and morphological results for dinemasporium-like and tiarosporella-like species do not correspond and this poses a taxonomic challenge. Thus, it has been suggested to use more gene regions for delineation at both generic and species level to have a better taxonomy result (Li et al. 2020). This also demonstrate that both morphological data and phylogenetic analyses are important to provide a more natural classification of coelomycetes (Li et al. 2020).

Molecular data availability has helped to link asexual taxa with their sexual morph (Shenoy et al. 2006, 2010, Li et al. 2020). Furthermore, under Article 59.1 of the International Code of Nomenclature for algae, fungi and plants, a single name is given for one fungus whether it is the asexual or sexual morph, known as “one fungus, one name” (Hawksworth 2012). Shenoy et al. (2006) proposed that for fungi that where sexual morph has not been detected, the asexual names should be used as holomorphs. Recently, Li et al. (2020) have provided a classification on hyaline coelomycetes, Wijayawardene et al. (2016) worked on the classification and taxonomy of dematiaceous coelomycetes, and Wijayawardene et al. (2020) provided a checklist for genera of coelomycetes.

Requirement for coelomycetes webpage

Several genera of coelomycetes lack molecular data or the holotypes are not available and this hampers their integration into a proper classification scheme. Mycologists usually compile all fungal work that are being carried out and publish them in the form of introduction and revision of

different taxa within each respective fungal class (Wijayawardene et al. 2016, 2017, Li et al. 2020, Wijayawardene et al. 2020, Hyde et al. 2020). Nevertheless, the lack of phylogenetic data and cultures complicate the understanding of the taxonomic concept of coelomycetes and make generic delimitation difficult (Li et al. 2020).

For the time being, it is difficult to gather taxonomic data on many coelomycetes as there is no specific platform to obtain information that is related only to coelomycetes. Data on coelomycetes are difficult to retrieve and are scattered. An electronic database makes information more accessible, user- and eco-friendly. Mycologist will be a few clicks away from discovering the ongoing work that is carried out on coelomycetes. Several websites have been created to help researchers to obtain mycological research work on one platform as mentioned in Table 1 and these updated on a regular basis (Jayasiri et al. 2015, Monkai et al. 2019, Pem et al. 2019a, b, Jayawardena et al. 2019, Bundhun et al. 2020).

The coelomycetes website, <https://coelomycetes.org/>, consists of only asexual fungi and (1) provides an up-to-date outline of coelomycetes; (2) provides notes on orders, families, genera and species of coelomycetes; and (3) provides updated taxonomic accounts of each genus that is in line with current research. The webpage also has a list of references that are linked to coelomycetes.

Table 1 List of websites

Website names	URL links	References
Marine fungi	https://marinefungi.org/	Jones et al. (2019)
Freshwater Fungi	https://freshwaterfungi.org/	Calabon et al. (2020)
One Stop Shop	https://onestopshopfungi.org/	Jayawardena et al. (2019)
Genera of Fungi	https://fungalgenera.org/	Monkai et al. (2019)
Dothideomycetes	https://dothideomycetes.org/	Pem et al. (2019a,b)
Sordariomycetes	https://sordariomycetes.org/	Bundhun et al. (2020)
Faces of Fungi	http://www.facesoffungi.org	Jayasiri et al. (2015)

Coelomycetes website

The website consists of up to-date taxonomic information on coelomycetes. Notes for each genus and species along with their taxonomic placement based on current DNA sequence data analyses (phylogenies) will be uploaded regularly. Each genus or species entry will have a description, photographic plate, phylogenetic tree, lists of accepted species in each genus, and include significant information through the links from other associated webpages (Table 1). Mycologists are invited to help improve and provide feedback on the webpage by checking the information that will be displayed. This webpage will definitely provide a better understanding of coelomycetes.

Construction

The coelomycetes website will include asexual fungi from recent classifications (Pem et al. 2019b, Hyde et al. 2020, Li et al. 2020, Bundhun et al. 2020). Index Fungorum and Faces of Fungi numbers for all species will be provided. Mycobank and herbaria numbers will also be given where available. Previously published data will be uploaded as new information are being published. The list of curators is enlisted in Table 2.

Table 2 List of curators.

Position	Name	Contact detail
Head Curators	Wijayawardene NN Li WJ	nalinwijayawardene@yahoo.com winnie20070653026@163.com
Managing Curator	Huanraluek N	naruemon.hua@mfu.ac.th

Table 2 Continued.

Position	Name	Contact detail
Senior curator	Jayawardena RS	ruvi.jaya@yahoo.com
Curators list	Bhunjun CS	avnishbhunjun@gmail.com
	Calabon MS	mscalabon@up.edu.ph
	Goonasekara ID	ishanidg@gmail.com
	Hongsanan S	sinang333@gmail.com
	Hyde KD	kdhyde3@gmail.com
	Jeewon R	r.jeewon@uom.ac.mu
	Maharachchikumbura SSN	sajeewa83@yahoo.com
	Norphanphoun C	oomchn@gmail.com
	Pem D	pem.dhandevi@gmail.com
	Phukhamsakda C	chayanard91@gmail.com
	Senanayake IC	indunilchinthani@gmail.com
	Shivas RG	roger.shivas@usq.du.au

Database interface and visualization

The website <https://coelomycetes.org/> is user-friendly. It helps with quick and easy research on coelomycetous taxa. The webpage consists of the following sections:

Home (Fig. 1)

The homepage provides an overview of the website, the objectives and “Highlights of Information”.

Outline (Figs 2, 3, 4)

The outline consists of recent taxonomic classification of species, genera, families, orders and subclass in coelomycetes.

Archives (Fig. 5)

The Archives provides consistent information on a specific order, family and genus.

Curators

This segment provides the list and contacts of people who are in charge of the webpage.

History

This section provides a brief history of coelomycetes.

References

This section provides references used in the text and are hyperlinked to facilitate their search online.

Notes

This section provides taxonomic information and any recent changes that occur on a specific coelomycete (Fig. 6; Recent genera and species uploaded).

Contact

The ‘Contact’ section helps to establish communication between users and curators. The users are allowed to add any comments and suggestions.




Fig. 1 – Homepage that provides an overview of the website, the objectives and “Highlights of Information”.




Fig. 2 – Outline consists of recent taxonomic classification of species, genera, families, orders and subclass in coelomycetes.

Phylum	ASCOMYCOTA, genera incertae sedis
	<i>Abrothallus</i> De Not.
	<i>Agryllopsis</i> Höhn.
	<i>Ajekarella</i> Kamat & Kalani.
	<i>Ajekarella polychaetriae</i> Kamat & Kalani, Mycopath.
	<i>Amerosporiopsis</i> Petr.
	<i>Anaphysmene</i> Bubák.
	<i>Angiopomopsis</i> Höhn.
	<i>Anthracoaderma</i> Speg.
	<i>Aoria</i> Cif.
	<i>Aphanofax</i> B. Sutton.
	<i>Aplocarpella</i> Syd & P. Syd.
	<i>Aristastoma</i> Tehon.
	<i>Ascochytopsis</i> Henn.
	<i>Ascochytilina</i> Petr.
	<i>Asteroconium</i> Syd & P. Syd.
	<i>Asteromidium</i> Speg.
	<i>Avettaea</i> Petr. & Syd.
	<i>Barnettella</i> D. Rao & P.Rag. Rao.
	<i>Bellificauda</i> B. Sutton.
	<i>Blennoria</i> Moug. & Fr.
	<i>Botryocrea</i> Petr.
	<i>Brencklea</i> Petrak.
	<i>Brycekendrickia</i> Nag Raj.
	<i>Brycekendrickia indica</i> Nag Raj, Can. J. Bot.
	<i>Caeruleoconidia</i> Zhurb & Diederich.
	<i>Callistospore</i> Petr.
	<i>Camarosporium</i> Tassi.
	<i>Camarosporiopsis</i> Abbas et al.
	<i>Capitrostrum</i> Bat.
	<i>Carnegieispora</i> Elayo & F. Berger.
	<i>Ceratopycnis</i> Höhn.
	<i>Chaetodiplodia</i> P. Karst.
	<i>Chithramia</i> Nag Raj.
	<i>Coniambigua</i> Elayo & Diederich.
	<i>Calocline</i> Syd.
	<i>Camatopycnis</i> E.K. Cash.
	<i>Catenophora</i> Luttr.
	<i>Ceuthodiplospora</i> Dled.
	<i>Chaetoseptoria</i> Tehon.
	<i>Cheilaria</i> Lib.
	<i>Choanatiara</i> DiCosmo.
	<i>Choanatiara gravis</i> Nag Raj, in Nag Raj & DiCosmo, Can. J. Bot.
	<i>Choanatiara lunata</i> DiCosmo & Nag Raj, in Nag Raj & DiCosmo, Can. J. Bot.
	<i>Chondropodiella</i> Höhn.
	<i>Ciliochora</i> Höhn.
	<i>Clypeopycnis</i> Petr.
	<i>Collonaemella</i> Höhn.
	<i>Comatospore</i> Piroz & Shoemaker.
	<i>Comatospore suttoni</i> Piroz. & Shoemaker, Can.
	<i>Cornucopella</i> Höhn.
	<i>Cornutispora</i> Piroz.
	<i>Cornutispora limaciformis</i> Piroz.
	<i>Crandallia</i> Ellis & Sacc.
	<i>Cryptomycella</i> Höhn.

Fig. 3 – Outline consists of recent taxonomic classification of species, genera, families, orders and subclass in coelomycetes (Continued).

Phylum	BASIDIOMYCOTA R. T. Moore.
Subphylum	AGARICOMYCOTINA Doweld.
Class	Agaricomycetes Doweld.
Subclass	Agaricomycetidae Parmasto.
Order	Agaricales underw.
Family	Phyalacriaceae Corner.
	<i>Mycotribulus</i> Nag Raj & W.B. Kendr.
	<i>Mycotribulus mirabilis</i> Nag Raj & W.B. Kendr.
Subclass	incertae sedis
Order	Corticiales K.H. Larss.
Family	Corticaceae Herter.
	<i>Giulia</i> Tassi.
	<i>Giulia tenuis</i> (Sacc.) Tassi ex Sacc. & D. Sacc.
Order	Sebacinales M. Weiss, Selosse, Rexer, A. Urb. & Oberw.
Order	Sebacinales, genus incertae sedis
	<i>Chaetospermum</i> Sacc.
	<i>Chaetospermum artocarp</i> (Nag Raj) Nag Raj.
	<i>Chaetospermum camelliae</i> Agnihottr.
Class	Agaricomycetes, genera incertae sedis
	<i>Cenangiomycetes</i> Dyko & B. Sutton.
	<i>Ellula</i> Nag Raj.
	<i>Fibulocoela</i> Nag Raj.
	<i>Fibulocoela indica</i> Nag Raj.
	<i>Pycnovellomyces</i> R.F. Castañeda.

Fig. 4 – Outline consists of recent taxonomic classification of species, genera, families, orders and subclass in coelomycetes (Continued).

COELOMYCETES
ISSN *****

Home Outline Archives Curators History References Notes Contact

Coelomyces Heirarchy

Agaricomycetes incertae sedis
[Read more about Agaricomycetes incertae sedis orders »](#)

Agaricomycetes incertae sedis

Ascomycota, genera

Basidiomycota, genera incertae sedis

Botryosphaeriales

Capnodiales

Capnodiales

Chaetomiales

Chaetosphaeriales

Conidiocarpus

Diaporthales

Glomerellales

Helotiales

Hypocreales

Medeolariales

Ostropales

Phacidiales

Phyllachorales

Pleosporales

Pleosporales

Rhizomatales

Sordariomycetes

Xylariales

Search Go

Recent News
[coelomyces.org - notes 2](#)
[coelomyces.org - notes 1](#)

Recent Genus
[Fuckella](#)
[Fibulocoela](#)
[Eutiarosporella](#)

Recent Species
[Fibulocoela indica](#)
[Eutiarosporella tritici](#)
[Eriosporella calami](#)

About Coelomyces

The website Coelomyces.org provides an up-to-date classification and account of all genera of the class Coelomyces.

Contact

Email:
coelomyces@gmail.com

Address:
 Mushroom Research Foundation, Chiang Rai
 57100, Thailand

Copyright © 2021 Coelomyces. All Rights Reserved.

Fig. 5 – Archives provides consistent information on a specific order, family and genus.

COELOMYCETES
ISSN *****

Home Outline Archives Curators History References Notes Contact

Coelomyces.org is a website dedicated to the taxonomy and classification of Coelomyces. The website focuses on providing an up-to-date outline of the Coelomyces with notes on orders, families, genera and species.

When using this website please cite: Li et al. 2020, Wijayawardene et al. 2016.

Search Go

Recent News
[coelomyces.org - notes 2](#)
[coelomyces.org - notes 1](#)

Recent Genus
[Fuckella](#)
[Fibulocoela](#)
[Eutiarosporella](#)

Recent Species
[Fibulocoela indica](#)
[Eutiarosporella tritici](#)
[Eriosporella calami](#)

Highlights of Information

- Provide illustrations and plates of Coelomyces genera

The website [Coelomyces.org](#) is updated periodically.

About Coelomyces **Contact**

Fig. 6 – Recent genera and species uploaded.

Acknowledgements

Authors are thankful to the Mushroom Research Foundation (MRF), Thailand. MS Calabon is grateful to Department of Science and Technology – Science Education Institute (Philippines).

References

- Bundhun D, Maharachchikumbura SSN, Jeewon R, Senanayake IC et al. 2020 – <https://sordariomycetes.org/>, a platform for the identification, ranking and classification of taxa within Sordariomycetes. *Asian Journal of Mycology* 3, 13–21. Doi 10.5943/ajom/3/1/2.
- Calabon MS, Hyde KD, Jones EBG, Chandrasiri S et al. 2020 – www.freshwaterfungi.org, an online platform for the taxonomic classification of freshwater fungi. *Asian Journal of Mycology* 3, 419–445. Doi 10.5943/ajom/3/1/14.
- Grove WB. 1919 – Mycological notes IV. *The London Journal of Botany* 57, 206–210.
- Gruyter JD, Woudenberg JHC, Aveskamp MM, Verkley GJM et al. 2017 – Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenochaeta* and *Pleurophoma*. *Mycologia* 102, 1066–1081.
- Hawksworth DL. 2012 – Managing and coping with names of pleomorphic fungi in a period of transition. *The International Mycological Association Fungus* 3, 15–24.
- Hughes SJ. 1953 – Conidiophores, conidia and classification. *Canadian Journal of Botany* 31, 577–659.
- Hyde KD, Norphanphoun C, Maharachchikumbura SSN, Bhat DJ et al. 2020 – Refined families of Sordariomycetes. *Mycosphere* 11, 305–1059.
- Li WJ, McKenzie E, Liu J, Bhat D et al. 2020 – Taxonomy and phylogeny of hyaline-spored coelomycetes. *Fungal Diversity* 100, 279–801.
- Liu F, Bonthond G, Groenewald JZ, Cai L et al. 2019 – *Sporocadaceae*, a family of coelomycetous fungi with appendage-bearing conidia. *Studies in Mycology* 92, 287–415.
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat DJ et al. 2015 – The Faces of Fungi database: fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74, 3–18.
- Jayawardena RS, Hyde KD, McKenzie EHC, Jeewon R et al. 2019 – One stop shop III: taxonomic update with molecular phylogeny for important phytopathogenic genera: 51–75 (2019). *Fungal Diversity* 98, 77–160.
- Jones EBG, Pang KL, Abdel-Wahab MA, Scholz B et al. 2019 – An online resource for marine fungi. *Fungal Diversity* 96, 347–433.
- Kendrick WB. 2000– *The Fifth Kingdom*, 3rd edn. Focus Publishing, Newbury.
- Monkai J, McKenzie EHC, Phillips AJL, Hongsanan S et al. 2019 – <https://fungalgenera.org/>: A web-based utility of classification and notes for all genera of fungi. *Asian Journal of Mycology* 2, 297–304.
- Nag Raj TR. 1981– *Coelomycete systematics Biology of conidial fungi* (Cole GT, Kendrick B), Academic Press, New York, USA, 43–84.
- Nag Raj TR. 1993– *Coelomycetous anamorphs with appendage-bearing conidia*. Mycologue Publications, Waterloo, 1–1101.
- Pem D, Jeewon R, Bhat DJ, Doilom M et al. 2019a – *Mycosphere Notes* 275–324: A morphotaxonomic revision and typification of obscure Dothideomycetes genera (incertae sedis). *Mycosphere* 10, 1115–1246. Doi 10.5943/mycosphere/10/1/22.
- Pem D, Hongsanan S, Doilom M, Tibpromma S et al. 2019b – <https://www.dothideomycetes.org>: An online taxonomic resource for the classification, identification, and nomenclature of Dothideomycetes. *Asian Journal of Mycology* 2(1), 287–297. Doi 10.5943/ajom/2/1/19.
- Shenoy BD, Jeewon R, Wang HK, KH Amandeep et al. 2010 – Sequence data reveals phylogenetic affinities of fungal anamorphs *Bahusutrabeeja*, *Diplococcium*, *Natarajania*, *Paliphora*, *Polyschema*, *Rattania* and *Spadicoides*. *Fungal Divers* 44, 161–169.

- Shenoy BD, Jeewon R, Wu WP, Bhat DJ, Hyde KD 2006 – Ribosomal and RPB2 DNA sequence analyses suggest that *Sporidesmium* and morphologically similar genera are polyphyletic. *Mycological Research* 110, 916–928.
- Sutton BC. 1980 – The Coelomycetes. Fungi imperfecti with pycnidia, acervuli and stromata. Commonwealth Mycological Institute, Kew, 1–696.
- Wijayawardene NN, Dai DQ, Tian Y, Tang LZ et al. 2020 – The genera of Coelomycetes, including genera of lichen forming, sexual morphs and synasexual morphs with coelomycetous morphs (genera A–C). *MycoAsia* 1–92.
- Wijayawardene NN, Hyde KD, Tibpromma S, Wanasinghe DN et al. 2017 – Towards incorporating asexual fungi in a natural classification: checklist and notes 2012– 2016. *Mycosphere* 8(9), 1457–1555, Doi 10.5943/mycosphere/8/9/10
- Wijayawardene NN, Hyde KD, Wanasinghe DN, Papizadeh M et al. 2016 – Taxonomy and phylogeny of dematiaceous coelomycetes. *Fungal Divers* 77, 1–316.
- Zhang SN, Hyde KD, Jones EBG, Jeewon R et al. 2019 – *Striatiguttulaceae*, a new pleosporalean family to accommodate *Longicorpus* and *Striatiguttula* gen. nov. from palms. *MycKeys* 49, 99–129.